

CEREZYME (IMIGLUCERASE)

INFUSION ORDERS

FAX: 470.922.3656 | **PHONE:** 470.377.6400

PATIENT INFORMATION: Fax completed form, insurance information, and clinical documentation to 470.922.3656		
Patient Name: DOB: Phone:		
Patient Status: ☐ New to Therapy ☐ Continuing Therapy Next Treatment Date:		
MEDICAL INFORMATION		
Diagnosis: ☐ Gaucher Disease ☐ Type 1 ☐ Type 3		
□ Type 1 □ Type 3 ICD-10 Code: E75.22		
10B-10 Gode. L75.22		
Patient Weight:Ibs. (required) Allergies:		
THERAPY ORDER		
Cerezyme: ☐ Dose: 60units/kg IV every 2 weeks x1 year		
□Other Dosage:		
Pre-Medication Orders: ☐ Tylenol 1000mg PO		
☐Benadryl 25 mg PO		
□Solumedrol mg IV		
□ Other: mg //		
Prescriber to monitor for antibody formation during 1st year of treatment.		
Lab Orders: Lab Frequency:		
Required labs to be drawn by:		
Required labs to be drawn by. \square blocare initiation \square Referring Frovider		
Required labs to be drawn by. Diocare initiation Referring Frovider		
Other orders:		
Other orders:		
Other orders: Anaphylactic Reaction Orders:		
Other orders: Anaphylactic Reaction Orders: Epinephrine (based on patient weight)		
Other orders: Anaphylactic Reaction Orders:		
Other orders: Anaphylactic Reaction Orders: Epinephrine (based on patient weight) 30kg (>66lbs): EpiPen 0.3mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x1		
Other orders: Anaphylactic Reaction Orders: Epinephrine (based on patient weight) 30kg (>66lbs): EpiPen 0.3mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x1 15-30kg (33-66lbs): EpiPen Jr. 0.15mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x1 Diphenhydramine: Administer 25-50mg orally OR IV (adult) Refer to physician order or institutional protocol for pediatric dosing		
Other orders: Anaphylactic Reaction Orders: Epinephrine (based on patient weight) 30kg (>66lbs): EpiPen 0.3mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x1 15-30kg (33-66lbs): EpiPen Jr. 0.15mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x1 Diphenhydramine: Administer 25-50mg orally OR IV (adult) Refer to physician order or institutional protocol for pediatric dosing Flush orders: NS 1-20mL pre/post infusion PRN and Heparin 10U/mL or 100U/mL per protocol as indicated PRN		
Other orders:		
Other orders: Anaphylactic Reaction Orders: Epinephrine (based on patient weight) 30kg (>66lbs): EpiPen 0.3mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x1 15-30kg (33-66lbs): EpiPen Jr. 0.15mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x1 Diphenhydramine: Administer 25-50mg orally OR IV (adult) Refer to physician order or institutional protocol for pediatric dosing Flush orders: NS 1-20mL pre/post infusion PRN and Heparin 10U/mL or 100U/mL per protocol as indicated PRN		
Other orders: Anaphylactic Reaction Orders: Epinephrine (based on patient weight) > 30kg (>66lbs): EpiPen 0.3mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x1 15-30kg (33-66lbs): EpiPen Jr. 0.15mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x1 Diphenhydramine: Administer 25-50mg orally OR IV (adult) Refer to physician order or institutional protocol for pediatric dosing Flush orders: NS 1-20mL pre/post infusion PRN and Heparin 10U/mL or 100U/mL per protocol as indicated PRN PROVIDER INFORMATION By signing this form and utilizing our services, you are authorizing Biocare Infusion, and its employees to serve as your prior authorization and specialty pharmacy designated agent in dealing with medical and prescription insurance companies, and to select the preferred site of care for the patient		
Other orders:		
Anaphylactic Reaction Orders: Epinephrine (based on patient weight) Solving (>66lbs): EpiPen 0.3mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x1 The solving (33-66lbs): EpiPen Jr. 0.15mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x1 Diphenhydramine: Administer 25-50mg orally OR IV (adult) Refer to physician order or institutional protocol for pediatric dosing Flush orders: NS 1-20mL pre/post infusion PRN and Heparin 10U/mL or 100U/mL per protocol as indicated PRN PROVIDER INFORMATION By signing this form and utilizing our services, you are authorizing Biocare Infusion, and its employees to serve as your prior authorization and specialty pharmacy designated agent in dealing with medical and prescription insurance companies, and to select the preferred site of care for the patient Provider Name: Signature: Date: Provider NPI: Phone: Fax: Contact Person: Opt out of Biocare Infusion selecting site of care (if checked, please list site of care):		
Other orders:		
Anaphylactic Reaction Orders: Epinephrine (based on patient weight) Solving (>66lbs): EpiPen 0.3mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x1 The solving (33-66lbs): EpiPen Jr. 0.15mg or compounded syringe IM or subQ; may repeat in 5-10 minutes x1 Diphenhydramine: Administer 25-50mg orally OR IV (adult) Refer to physician order or institutional protocol for pediatric dosing Flush orders: NS 1-20mL pre/post infusion PRN and Heparin 10U/mL or 100U/mL per protocol as indicated PRN PROVIDER INFORMATION By signing this form and utilizing our services, you are authorizing Biocare Infusion, and its employees to serve as your prior authorization and specialty pharmacy designated agent in dealing with medical and prescription insurance companies, and to select the preferred site of care for the patient Provider Name: Signature: Date: Provider NPI: Phone: Fax: Contact Person: Opt out of Biocare Infusion selecting site of care (if checked, please list site of care):		



COMPREHENSIVE SUPPORT FOR

CEREZYME (IMIGLUCERASE) THERAPY

PATIENT INFORMATION:	
Patient Name:	DOB:
REQUIRED DOCUMENTATION FOR REFERRAL PROCESSING & INSURANCE APPROVAL	
☐ Include <u>signed and completed order</u> (MD/prescriber to complete page 1)	
☐ Include patient demographic information and insurance information	
☐ Include patient's medication list	
Supporting clinical notes to include any past tried and/or failed therapies, intolerance, benefits, or contraindications to conventional therapy	
 □ Does the patient have symptomatic Gaucher severe anemia, thrombocytopenia, bone disesplenomegaly? □ Yes □ No 	· ·
Does the patient have a history of failure or	intolerance to VPRIV? ☐ Yes ☐ No
☐ Include labs and/or test results to support diagr	nosis
☐ CBC, Hepatic Function Tests	
Other medical necessity:	

Biocare Infusion will complete insurance verification and submit all required documentation for approval to the patient's insurance company for eligibility. Our team will notify you if any additional information is required. We will review financial responsibility with the patient and refer him/her to any available co-pay assistance as needed. Thank you for the referral.

Please fax all information to (470) 922-3656 or call (470) 377-6400 for assistance